D-21358

## IN THE CLAIMS

- 1. (Currently amended) A one pot method for large scale production of an organometallic compound comprising a transition metal-containing amide, a transition metal-containing alkoxide, a transition metal-containing diketonate or a transition metal-containing imide, said one pot method comprising (i) reacting a hydrocarbon or heteroatom-containing material with a base material in the presence of a solvent and under reaction conditions sufficient to produce a first reaction mixture comprising a hydrocarbon or heteroatom-containing compound, (ii) adding a transition metal source compound to said first reaction mixture, (iii) reacting said hydrocarbon or heteroatom-containing compound with said transition metal source compound under reaction conditions sufficient to produce a second reaction mixture comprising said organometallic compound, and (iv) separating said organometallic compound from said second reaction mixture.
- (Currently amended) The one pot method of claim 1 wherein the large scale production amounts to about 0.25 kilograms or greater of said organometallic compound.
- (Currently amended) The one pot method of claim 1 wherein the large scale production amounts to about 0.5 kilograms or greater of said organometallic compound.
- 4. (Currently amended) The one pot method of claim 1 wherein the large scale production amounts to about 1.0 kilograms or greater of said organometallic compound.

+2038372515

FROM-Praxair, Inc.

- (Currently amended) The one pot method of claim 1 wherein the 5. organometallic compound yield is from about 75 to 99%-or greater.
- (Currently amended) The one pot method of claim 1 wherein the 6. organometallic compound yield is from about 80 to 99%-or greater.
- (Currently amended) The one pot method of claim 1 wherein the 7. hydrocarbon or heteroatom-containing material comprises an amine, alcohol, diketone, eyelepentadiene, imine, hydrocarbon or halogen.
- (Original) The one pot method of claim 1 wherein the base material 8. has a pKa greater than about 10.
- (Original) The one pot method of claim 1 wherein the base material 9. comprises BuLi, MeLi, NaH, CaH or a lithium amide.
- (Currently amended) The one pot method of claim 1 wherein the 10. hydrocarbon or heteroatom-containing compound comprises a lithiated amide, a lithiated alkoxide, a lithiated diketonate, eyelopentadienide or a lithiated imide.
- (Currently amended) The one pot method of claim 1 wherein the 11.  $\underline{transition}\ metal\ source\ compound\ is\ represented\ by\ the\ formula\ MX_n\ wherein\ M\ is\ a$ transition metal, X is halide and n is a value of 3, 4 or 5.
- (Original) The one pot method of claim 11 wherein the transition 12. metal is selected from hafnium, titanium, zirconium, tantalum and molybdenum.

D-21358

- 13. (Currently amended) The one pot method of claim 1 wherein the transition metal source compound comprises HfCl<sub>4</sub>, HfF<sub>4</sub>, HfBr<sub>4</sub>, HfI<sub>4</sub> or Hf(OTf)<sub>4</sub>.
- 14. (Currently amended) The one pot method of claim 1 wherein the transition metal source compound is HfCl4.
- 15. (Currently amended) The one pot method of claim 1 wherein the transition metal source compound is added to the first reaction mixture at ambient temperature or at a temperature greater than ambient temperature.
- 16. (Original) The one pot method of claim 1 wherein the solvent is selected from saturated and unsaturated hydrocarbons, aromatic hydrocarbons, aromatic heterocycles, alkyl halides, silylated hydrocarbons, ethers, polyethers, thioethers, esters, thioesters, lactones, amides, amines, polyamines, nitriles, silicone oils, other aprotic solvents, or mixtures of one or more of the above.
- 17. (Original) The one pot method of claim 1 wherein the solvent is selected from hexanes, THF or mixtures thereof.
  - 18. (Canceled)
- 19. (Original) The one pot method of claim 1 wherein the organometallic compound comprises hafnium amide, hafnium (IV) tert-butoxide, hafnium (IV) acetylacetonate, bis(cyclopentadienyl)hafnium dichloride or t-butylimidobis(dimethylamino)hafnium.
- 20. (Currently amended) A one pot method for producing a liquid-hafnium amide compound comprising (i) reacting an amine with a lithiated base material to

09:10

D-21358

produce a first reaction mixture comprising a lithium amide, (ii) adding a hafnium halide to said first reaction mixture, (iii) reacting said lithium amide with said hafnium halide under reaction conditions sufficient to produce a second reaction mixture comprising said liquid-hafnium amide compound, and (iv) separating said liquid-hafnium amide compound from said second reaction mixture.

- organometallic compound comprising a transition metal-containing amide, a transition metal-containing alkoxide, a transition metal-containing diketonate, a transition metal-containing cyclopentadienide or a transition metal-containing imide, said one pot method comprising (i) reacting a hydrocarbon or heteroatom-containing material with a base material in the presence of a solvent and under reaction conditions sufficient to produce a first reaction mixture comprising a hydrocarbon or heteroatom-containing compound, (ii) adding a transition metal source compound to said first reaction mixture, wherein said transition metal source compound is added to said first reaction mixture at ambient temperature or at a temperature greater than ambient temperature, (iii) reacting said hydrocarbon or heteroatom-containing compound with said transition metal source compound under reaction conditions sufficient to produce a second reaction mixture comprising said organometallic compound, and (iv) separating said organometallic compound from said second reaction mixture.
- 22. (New) A one pot method for large scale production of an organometallic compound comprising a transition metal-containing amide, a transition metal-containing alkoxide, a transition metal-containing diketonate, a transition metal-containing cyclopentadienide or a transition metal-containing imide, said one pot method comprising (i) reacting a hydrocarbon or heteroatom-containing material with a base material in the presence of a solvent and under reaction conditions sufficient to produce a first reaction mixture comprising a hydrocarbon or heteroatom-containing

D-21358

compound, (ii) adding a transition metal source compound to said first reaction mixture, (iii) reacting said hydrocarbon or heteroatom-containing compound with said transition metal source compound under reaction conditions sufficient to produce a second reaction mixture comprising said organometallic compound, and (iv) separating said organometallic compound from said second reaction mixture; wherein the organometallic compound yield is from about 80 to 99%.

organometallic compound comprising (i) reacting a hydrocarbon or heteroatom-containing material with a base material in the presence of a solvent and under reaction conditions sufficient to produce a first reaction mixture comprising a hydrocarbon or heteroatom-containing compound, wherein said solvent is selected from hexanes, THF or mixtures thereof, (ii) adding a metal source compound to said first reaction mixture, (iii) reacting said hydrocarbon or heteroatom-containing compound with said metal source compound under reaction conditions sufficient to produce a second reaction mixture comprising said organometallic compound, and (iv) separating said organometallic compound from said second reaction mixture.